RESEARCH ARTICLE

Exploring the structural characteristics of the Adult Social Care Outcomes Toolkit (ASCOT) and ASCOT-Carer [version 1; peer review: 2 approved with reservations]

Stacey Rand ¹, Ann-Marie Towers ², Juliette Malley³, Barbora Silarova¹

¹Personal Social Services Research Unit, University of Kent, Canterbury, UK

²Centre for Health Services Studies, University of Kent, Canterbury, UK

³Care Policy and Evaluation Centre, London School of Economics and Political Science, London, UK

V1 First published: 24 Feb 2022, 2:21
<https://doi.org/10.3310/nihropenres.13259.1>

Latest published: 21 Mar 2023, 2:21
<https://doi.org/10.3310/nihropenres.13259.2>

Abstract

Background: Measurement models inform the approach to assess a measure's validity and also how a measure is understood, applied and interpreted. With preference-based measures, it is generally accepted that they are *formative*; however, if they are applied without preferences, they may be *reflective*, *formative* or *mixed*. In this study, we sought to empirically test whether the *reflective*, *formative* or *mixed* measurement model best describes PBMs of social care-related quality of life (ASCOT, ASCOT-Carer). We also explored the network approach, as an alternative.

Methods: ASCOT and ASCOT-Carer data were analyzed using confirmatory factor analysis and Multiple Indicators Multiple Causes models to test reflective, formative or mixed measurement models, respectively. Network analysis of partial correlations using the Gaussian graphical model was also conducted.

Results: The results indicated that the reflective measurement model is the worst fit for ASCOT and ASCOT-Carer. The formative or mixed models may apply to ASCOT. The mixed model was the best fit for ASCOT-Carer. The network analysis indicated that the most important or influential items were *Occupation* and *Personal cleanliness and comfort* (ASCOT) and *Time and space* and *Self-care* (ASCOT-Carer).

Conclusions: The ASCOT and ASCOT-Carer are best described as formative/mixed or mixed models, respectively. These findings may guide the approach to the validation of cross-culturally adapted and translated versions. Specifically, we recommend that EFA be applied to establish structural characteristics, especially if the measure will be applied as a PBM and as a measure of SCRQoL. Network analysis may also provide further useful insights into structural characteristics.

Keywords

quality of life, social care, long-term care, ASCOT, service users, carers

Open Peer Review

Approval Status ? ?

1

2

version 2

(revision)

21 Mar 2023

version 1

24 Feb 2022

view

view

1. **You-Shan Feng** , Medical University of Tübingen, Tübingen, Germany

2. **Herdiantri Sufriyana** , Taipei Medical University, Taipei, Taiwan
 Universitas Nahdlatul Ulama Surabaya, Surabaya, Indonesia

Any reports and responses or comments on the article can be found at the end of the article.

Corresponding author: Stacey Rand (s.e.rand@kent.ac.uk)

Author roles: **Rand S:** Conceptualization, Formal Analysis, Funding Acquisition, Methodology, Project Administration, Supervision, Writing – Original Draft Preparation, Writing – Review & Editing; **Towers AM:** Conceptualization, Funding Acquisition, Writing – Review & Editing; **Malley J:** Conceptualization, Methodology, Writing – Review & Editing; **Silarova B:** Methodology, Writing – Review & Editing

Competing interests: No competing interests were disclosed.

Grant information: This research was funded by the National Institute for Health Research (NIHR) under its Research for Social Care (RfSC) Programme (Grant Reference Number: NIHR200058). The views expressed are those of the authors and not necessarily those of the NIHR or the Department of Health and Social Care.

The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Copyright: © 2022 Rand S *et al.* This is an open access article distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

How to cite this article: Rand S, Towers AM, Malley J and Silarova B. **Exploring the structural characteristics of the Adult Social Care Outcomes Toolkit (ASCOT) and ASCOT-Carer [version 1; peer review: 2 approved with reservations]** NIHR Open Research 2022, 2:21 <https://doi.org/10.3310/nihropenres.13259.1>

First published: 24 Feb 2022, 2:21 <https://doi.org/10.3310/nihropenres.13259.1>

Plain english summary

For many people living with long-term health conditions or disabilities, community-based social care services (like, home care) enable them to maintain independence, stay connected, and to live well. For families and friends who care for someone ('carers'), these services may also help them. They may allow carers to continue in paid employment and to have time for hobbies, friendships, to stay healthy, and connect with others in a similar situation. An important question is what type(s) of community-based services, best support people and their carers.

To find this out, we need a way of measuring the effect services have on people's lives. The Adult Social Care Outcomes Toolkit (ASCOT) is a questionnaire that asks people about aspects of their life that might be affected by social care services (for example, having control over everyday life). This questionnaire has already been used by researchers and care providers to review how well services support people. There is also another version of the questionnaire called the ASCOT-Carer, which looks at aspects of life that are important to carers.

There has been interest in culturally adapting and translating these measures into other languages. However, there are different ways of establishing how well a translated version relates back to the original. This is important to make sure that the new version is measuring what we expect it to.

In this paper, we compare different ways of understanding the information collected using the ASCOT and ASCOT-Carer in England. This will inform how to approach the testing of ASCOT (and other similar measures) that have been translated into new languages. It also helps us to understand how different aspects of life that are supported by social care services are related

to each other. This can inform our understanding of people's needs and how to best support them.

Introduction

The theoretical and philosophical questions of measurement models are important to psychometric research. Implicitly and explicitly, they inform the approach to assess a measure's validity and also how a measure is understood, applied and interpreted. In this paper, we will draw on a preference-based measure (PBM) used in economic evaluation of long-term care services, the Adult Social Care Outcomes Toolkit for service users (ASCOT)¹ and carers (ASCOT-Carer)^{2,3}, to illustrate the issues related to applying measurement models to PBMs. In doing so, we will highlight key implications for the development and application of PBMs in research, evaluation and practice.

The two measurement models commonly applied in psychometric research are *reflective measurement models* (RMM) and *formative models* (FM) (see [Figure 1](#)). In RMM, the construct is the common cause of items (*observables*), *i.e.* the relationships between the items are due to a common causal path to the construct, not interrelationships between the items. Compared to RMM, the items in FMs are conceptualized as a set of independent measures that come together to form a construct. The items may be inter-correlated. These models also differ in underlying philosophical assumptions. RMMs are based on a realist stance; the construct is conceptualized as being 'out there', but unobserved. By contrast, FMs are based on a constructivist position; the construct is a rational construction of the mind and is a theoretical composite of its constituent parts.

Typically, PBMs used in economic evaluation of health and social care services (also known as long-term care outside of the

Figure 1. Formative and Reflective Measurement Models.

UK), like the EQ-5D and ASCOT, are understood using the FM due to the composite nature of PBMs^{4,5}. Each item represents a dimension of health or care-related quality of life (QoL) that is distinct from the other items (the *observables*). The PBM index score represents an individual's outcome state (the *construct*). It is a composite function of the preference weights assigned to each item. The items for PBMs are usually selected on the basis that they will be weakly associated, to avoid redundancy and also to allow tradeoffs between items. Therefore, the proposed methodology for evaluating PBMs, based on the FM, focuses on content validity, face validity and construct validity of the descriptive system (without preferences), rather than structural validity⁵. The justification for this approach is that the classical test theory (CTT) method of establishing structural validity using exploratory factor analysis (EFA) is theoretically based on the RMM^{6,7}. It has been argued that these methods are irrelevant and potentially misleading for measures based on the FM^{8,9}, even if other CTT methods (e.g. confirmatory factor analysis) may be applied⁷.

However, PBMs, like ASCOT, may be used in a range of contexts. The ASCOT and ASCOT-Carer have been used as PBMs in economic evaluation^{10–12}; however, they have also been used in non-economic contexts, without preference weights, e.g. for needs assessment, care planning^{13,14}. Correspondingly, psychometric testing of these measures have taken a combination of RMM (i.e. using EFA)^{1,2} and FM (i.e. not applying structural validity methods) approaches^{15,16}, typically influenced by the view of how the measure will be applied (i.e. as a PBM, or not). Inconsistent approaches to psychometric evaluation of the same measure have given rise to discussion of the correct approach to the translation and cross-cultural validity or adaptation of the measures for new contexts or populations^{15–17}. One way of resolving this question is to establish empirically which measurement model, RMM or FM, most accurately describes the ASCOT and ASCOT-Carer.

In recent years, however, there have been advances in thinking about measurement models, beyond RMM and FMs. Instead, some measures may be best described by *mixed measurement models*, with a combination of formative and reflective relationships between construct and observables¹⁸. It has been proposed that measures based on mixed measurement models may be treated as RMMs (i.e. by applying EFA) in psychometric research⁸. However, there has been critique of limitations of RMM, FM and mixed models, which broadly apply also to the care-related QoL measures, ASCOT and ASCOT-Carer. For example, RMMs are based on the assumption that the observables are locally independent when controlling for the latent variable. This is unlikely to hold for ASCOT measures, since we expect there to be associations between observables (ASCOT or ASCOT-Carer items), aside from an underlying association with the latent construct (social care-related quality of life: SCRQoL), e.g., it is likely that a person's sense of control over their daily life would be directly affected by whether they feel they are doing things they value and enjoy; and *vice versa*. It is also conceptually implausible to say that the items have a common cause

(i.e. that having a poor SCRQoL will *result in* having poor control over daily life, personal safety, *etc.*), especially as ASCOT is a PBM where the items were selected to vary independently. Aside from general critique of FMs and their limitations, especially in application to psychological measurement (for example^{19–21}), a key limitation of FMs is that relationships between observables in FMs are modelled as noise. Where there is justification for expected relationships between observables, as outlined above for ASCOT, the formative approach potentially overlooks important structural information^{7,22}.

An alternative to RMMs, FMs or their combination in mixed models, is the *network model* (NM). NMs avoid the limitations of FMs, RMMs and mixed models by proposing instead that the construct is a network of causally-related elements (nodes), without any assumptions about the nature or causal direction of relationships⁷. NMs do not require the existence of latent variable(s), since the construct is still 'real' as a complex network or system of interrelated variables²³. The NM is based on a critical realist position, i.e. that the construct *is* the observed variables in their complex interrelationships of mutual influence²⁴. NMs have been applied in psychological measurement of intelligence²⁵, personality²⁶ and psychological comorbidity^{27,28} and has been used in health psychology research, as they enable modeling of complex interdependencies between factors that may affect an outcome²⁹. NMs have also been proposed as a method for psychometric analysis of health-related QoL measures, to address the limitations of both FM and RMMs when applied to measures of health-related QoL⁷.

The aim of this study was to establish which measurement model best describes the ASCOT and ASCOT-Carer, respectively: *reflective*, *formative* or *mixed*. While the development and psychometric assessment of the ASCOT instruments has applied either the FM¹⁶ or the RMM^{1,2}, these are directly compared here, alongside also mixed models. This will inform approaches to the future validation and development of ASCOT or related PBMs of care-related QoL, including in translation or cross-cultural adaptation. The network approach (NM) was also explored, as an alternative to RMMs, FMs and mixed models. The purpose was to establish whether it provides further insight into the structure of ASCOT and ASCOT-Carer, beyond what is offered by factor analysis, as a network of mutual interrelationship between items.

Methods

Sampling and data collection

To explore the internal structural characteristics of ASCOT and ASCOT-Carer, we conducted secondary analysis on data from two cross-sectional studies in England. These are described below. **Study One** collected data using the standard self-completion version (SCT4) of ASCOT^{1,30}. **Study Two** collected data from carers of people with dementia about their own QoL outcomes (ASCOT-Carer SCT4³). To reflect the specific needs and experiences of carers, the ASCOT-Carer has a different set of attributes to the service user versions, with some overlapping domains (see Box 1).

Box 1. SCRQoL attributes

ASCOT	ASCOT-Carer
Control over daily life	Control over daily life
Occupation (<i>doing things I value and enjoy</i>)	Occupation (<i>doing things I value and enjoy</i>)
Social participation and involvement	Social participation and involvement
Personal safety	Personal safety
Food and drink	
Accommodation comfort and cleanliness	
Personal comfort and cleanliness	
Dignity	
	Self-care (<i>being able to look after myself</i>)
	Time and space to be myself
	Feeling supported and encouraged in caring role

Study One: Identifying the Impact of Adult Social Care (IIASC)

The Identifying the Impact of Adult Social Care (IIASC) study was an interview survey of 990 users of community-based support/services in England. The survey was conducted in 22 local authorities (LAs) between June 2013 and March 2014. The sample was identified from records held by LAs or home care providers. The inclusion criteria were: aged 18 years or over, living in their own home, and receiving support due to physical disability or sensory impairment or mental health conditions or learning disabilities. Because the questionnaire for people with learning disabilities used an adapted easy read version of ASCOT, these data ($n=220$) are excluded from the analysis presented here.

Eligible participants were invited to participate in an interview, which was either completed face-to-face (74.2%) or by telephone (25.8% of the sample ($n=770$)). Written or verbal consent was obtained before all interviews. Data were collected on the respondent's personal characteristics, social care needs, health, type and intensity of service use, informal support from family/friends, and quality of life outcomes, including the ASCOT¹. Further details of the questionnaire content and data collection methods are outlined elsewhere³⁰.

Study Two: Measuring the Outcomes of People with Dementia and their Carers (MOPED) study

The MOPED study was an observational cross-sectional study to establish the psychometric properties of the ASCOT-Proxy and ASCOT-Carer. The data were collected using self-administered questionnaire (either postal questionnaire or an online version

in Qualtrics) among 313 unpaid family carers in England. The inclusion criteria were carers, who provided unpaid help or support to a relative, partner/spouse or friend living with dementia, who used community-based social care (*e.g.* home care, day centre), was not in residential or nursing care, and unable to self-complete a structured questionnaire, even with help.

Participants were recruited between January 2020 and April 2021 through *Join Dementia Research* (an online opt-in volunteer panel), local carers' support organisations, healthcare settings and social media. Written informed consent was obtained from all participants. The questionnaire collected data on the respondent's characteristics, caregiving situation and care-recipient characteristics. SCRQoL was measured by the ASCOT-Carer.

Statistical analysis

The first aim of the study was to empirically compare the three measurement models (*reflective*, *formative* and *mixed*) for ASCOT and ASCOT-Carer, to determine which fits best. The *reflective measurement model* was evaluated using confirmatory factor analysis. Each of the eight ASCOT and seven ASCOT-Carer items were tested separately as reflective indicators of a single latent construct (*i.e.* SCRQoL → ASCOT or ASCOT-Carer items – see Figure 2). As ASCOT and ASCOT-Carer items are categorical, not continuous, the models were estimated using weighted least squares.

The *formative* and *mixed models* were evaluated using Multiple Indicators Multiple Causes (MIMIC) models³¹. These are structural equation models (SEMs) that allow the simultaneous modelling of reflective items that relate to one or more latent variable(s), alongside the relationship between formative items and the latent variable. In the *formative model*, all eight or seven items of the ASCOT or ASCOT-Carer, respectively, were modelled with a formative relationship to the latent variable, SCRQoL. To enable the empirical testing of the models, it is necessary to specify also two or more reflective indicators. These are typically measures of the same or similar constructs. This provides external anchoring against validated measures or items (*i.e.* EQ-5D, QoL item), when all of the ASCOT items are considered as a composite of SCRQoL (*i.e.* ASCOT or ASCOT-Carer items → SCRQoL → EQ-5D, overall QoL – see Figure 3). Since there are no other validated measures of social care-related QoL than the ASCOT or ASCOT-Carer, measures of the related constructs of health-related QoL and overall QoL were selected. From previous research, these measures are known to be related to ASCOT SCRQoL^{2,12,32,33}. Specifically, for ASCOT, the EQ-5D-3L and a single item 7-point rating of overall QoL were considered. For ASCOT-Carer, the EQ-5D-5L converted by cross-walk to EQ-5D-3L values³⁴ and a single-item 5-point overall rating of QoL were considered. These differences in the measures were due to the available data in each study dataset.

In the *mixed models* (see Figure 4), the ASCOT or ASCOT-Carer items were considered as formative or reflective. These were selected based on theory. In the development of ASCOT, it has been noted that the domains may be categorized as those that relate to: (1) *basic domains* that relate to basic care

Figure 2. a. Reflective Model for ASCOT. b. Reflective Model for ASCOT-Carer.

Figure 3. a. Formative Model for ASCOT. b. Formative Model for ASCOT-Carer.

Figure 4. **a.** Mixed Model for ASCOT. **b.** Mixed Model for ASCOT-Carer. **Key to Figures 2–4:** Acc, accommodation; Pers, personal comfort and cleanliness; Food, food and drink; Safe, personal safety; Occ, occupation; Con, control over daily life; Soc, social participation; Dig, dignity; Self-care, self-care; Time, time and space to be yourself; Supp, feeling supported and encouraged in caring role.

needs/support to sustain life and health (*i.e.* Food and drink, Personal or Accommodation comfort and cleanliness, Personal safety (ASCOT); Self-care, Feeling supported in the caring role, Personal safety (ASCOT-Carer)); (2) *higher order domains* that relate to aspects of QoL beyond basic care needs and/or relate to a person's sense of self and identity (*i.e.* Control over daily life, Occupation, Social participation (ASCOT and ASCOT-Carer)); (3) domains related to how the delivery of care affects a person's sense of self and identity (Dignity (ASCOT) or Time and space to be yourself (ASCOT-Carer))³³. The *basic domains* were considered as formative, since they may be conceptualized as constituent parts of social care-related QoL (*i.e.* aspects of QoL that make up the construct, SCRQoL). The *higher order domains* and the *domains related to sense of self and identity* were considered as reflective of SCRQoL (*i.e.* they are driven by a common factor, SCRQoL).

To compare these models, standardized factor loadings and coefficients were reported for the CFA and MIMIC, respectively, to explore the relationship between items and the latent variable. Model fit statistics were calculated to evaluate the estimated models. The following criteria were applied to indicate good fit: root mean square error of approximation (RMSEA) ≤ 0.06 (upper confidence interval of ≤ 0.08), standardized root mean square residual (SMSR) of ≤ 0.08 , with a comparative fit index (CFI) and Tucker-Lewis index (TLI) of ≥ 0.95 ³⁵.

In addition to comparing and evaluating RMM, FM and mixed models, we also applied network analysis to determine whether this approach, which has been proposed as a suitable approach for measures of health-related QoL⁷, offers additional useful insights when applied to social care-related QoL measures, like ASCOT. In network analysis, the focus is on the variables

(**nodes**) and the relationships between them (**edges**). These may be **directed**, which indicate a one-way effect, or **undirected**, which indicates an unspecified mutual relationship.

In this analysis, the ASCOT and ASCOT-Carer items were considered as the **nodes** and all **edges** were specified as undirected. The network was estimated by analyzing partial correlations using the Gaussian graphical model. Polychoric correlation coefficients were applied since the data were ordinal. The coefficients are estimates of the strength of relationship between variables (the ASCOT or ASCOT-Carer items) whilst controlling for the effects of other measured variables in the model. A graphical least absolute shrinkage and selection operator (*lasso*) approach was applied to the estimation of the correlation network³⁶. This statistical technique takes into account the model complexity and seeks to reduce the number of spurious relationships by reducing small weak edge estimates to zero. The *lasso* tuning parameter (λ) may be set from 0 to 1. Increasing the tuning parameter will minimize spurious edges, however, relevant edges may also be suppressed²⁹. The analysis applied $\lambda=0.25$.

After the models were estimated, the network properties were evaluated. The centrality of nodes (*i.e.* their relative importance) in determining the network structure was assessed by the number of connections incident to the node (*degree centrality*). Centrality indices were also calculated and reported for node *strength*, *closeness* and *betweenness*²⁹. The strength index is a composite measure of both the number and strength of connections to a node. The closeness index represents the relationship between one node and the other nodes through its indirect connections (*i.e.* its connectedness or connectivity). A high closeness index indicates that the node is affected quickly by changes to the other nodes in the network. The betweenness

index indicates the importance of a node in relation to the average pathway between other nodes²⁹.

The descriptive statistics, CFA and MIMIC were calculated in STATA version 16. The network analysis was conducted in R.

Results

The sample characteristics are outlined in [Table 1](#). The sample for study 1 (IIASC) was users of social care services. Just over half the sample (52.7%) were aged 65 years or older and 42% were male. There was a wide range of social care needs in the sample, with 9.4% of respondents reporting no needs for activities of daily living. (The study sample included users of services for support with mental health difficulties, where the eight ADLs may not relate to the person's needs.)

The study 2 (MOPED) sample were all carers of someone with dementia. The majority of the MOPED sample (study 2) were caring for a parent (n=152, 48.6%) or a spouse or partner (n=130, 41.5%). Most carers were co-resident with the person they support (n=181, 57.8%). The high level of social care need of care recipients is reflected in the profile of difficulty with ADLs; 32.2% of the sample reported that the person they supported had difficulty with all eight ADLs. Almost half of the sample (47.0%) were carers providing 50 or more hours of unpaid care per week.

The distribution of scores by item for ASCOT (IIASC, Study 1) and ASCOT-Carer (MOPED, Study 2) are shown in [Figure 5](#) and [Figure 6](#). The rating of high-level or some needs are highest for the three ASCOT 'higher order' domains of Control over

Table 1. Sample characteristics.

	Study 1 (n=770)	Study 2 (n= 313)
	N (%)	N (%)
Male	323 (42.0%)	76 (24.3%)
Age ≥65 years	406 (52.7%)	137 (43.8%)
Ethnicity: white British	704 (91.4%)	296 (94.6%)
Self-reported health: <i>good or very good</i>	228 (29.6%)	229 (73.2%)
<i>fair</i>	311 (40.4%)	72 (23.0%)
<i>bad or very bad</i>	230 (29.9%)	11 (3.5%)
<i>missing data</i>	1 (0.1%)	1 (0.3%)
Care recipient's ADLs with difficulty ¹ :		
<i>none</i>	72 (9.4%)	0 (0%)
<i>1–4</i>	293 (38.1%)	123 (39.3%)
<i>5–7</i>	243 (31.6%)	86 (27.5%)
<i>All 8</i>	159 (20.6%)	101 (32.2%)
<i>missing data</i>	3 (0.4%)	3 (1.0%)
Hours of inf care / week: 0–19 hours	n/a	95 (30.4%)
20–34 hours	n/a	36 (11.5%)
35–50 hours	n/a	32 (10.1%)
≥50 hours	n/a	147 (47.0%)
<i>missing data</i>	n/a	3 (1.0%)
	Mean (Std. Dev., Range)	Mean (Std. Dev., Range)
Overall QoL	4.43 (1.26, 1 to 7)	3.46 (1.04, 1 to 5)
EQ-5D-3L Index	0.27 (0.39, –0.594 to 1)	0.79 (1.04, –0.594 to 1)

¹ This is a count of activities of daily living (ADLs) where the respondent (or care recipient by proxy report) had difficulty or was unable to complete the task alone, without help.

Figure 5. ASCOT ratings (Study 1).

Figure 6. ASCOT-Carer ratings (Study 2).

daily life, Social participation and Occupation. The ideal state (best care-related QoL) was rated by over half of the sample for

the basic domains of Personal comfort and cleanliness (56.1%), Accommodation (60.1%) and Food and drink (70.1%).

The ASCOT-Carer ratings had a different response profile. With the exception of Personal safety (3.2%), between 28% (Occupation) and 63% (Time and space) of ratings per attribute were high-level or some needs. This is broadly consistent with a 2013/14 data collection from carers in England using the ASCOT-Carer². However, the current study sample had a higher profile of need with regard to Control over daily life (51.8% *versus* 37.0%) and Social participation (58.8% *versus* 33.3%). The reverse was the case for Occupation (28.8% *versus* 49.1% of the sample reported high-level or some needs)². This is unsurprising given that the study sample was carers of people with dementia living in their own homes. This subgroup of carers are known to have specific high-level social care support needs that relate to the condition, for example, in its fluctuating and unpredictable

nature^{37–39}. The data collection also coincided with the COVID-19 pandemic and associated restrictions in England, which may have affected a number of aspects of quality of life due to the legal restrictions. The lower levels of QoL for Social participation and Control over daily life may, at least partly, be affected by the legal restrictions on socializing, travel/movement, leaving the home and other aspects of life designed to curb the spread of the infection.

The results of the reflective, formative and mixed models are shown in Table 2. The reflective models were a poor fit against the criteria (*i.e.* RMSEA ≤ 0.06 (upper confidence interval ≤ 0.08), SMSR ≤ 0.08 , and CFI/TLI ≥ 0.95 ³⁵). None of the criteria were met for ASCOT; only SMSR ≤ 0.08 was met for ASCOT-Carer.

Table 2. Confirmatory factor analysis (reflective model) and MIMIC (formative and mixed models).

	Confirmatory factor analysis (reflective)				MIMIC (formative)				MIMIC (mixed)			
	Study 1: ASCOT		Study 2: ASCOT-C		Study 1: ASCOT		Study 2: ASCOT-C		Study 1: ASCOT ¹		Study 2: ASCOT-C ²	
	Factor loading	SE	Factor loading	SE	Stand. Coeff.	SE	Stand. Coeff.	SE	Stand. Coeff.	SE	Stand. Coeff.	SE
Accommodation	0.388**	0.053			0.048	0.045			0.124**	0.043		
Food and drink	0.314**	0.049			0.089*	0.042			0.128**	0.040		
Personal care	0.461**	0.048			0.078	0.046			0.308**	0.041		
Personal safety	0.412**	0.043	0.334***	0.055	0.177**	0.042	0.133*	0.055	0.238**	0.039	0.094	0.052
Control	0.609**	0.033	0.747***	0.032	0.247**	0.046	0.205**	0.065	0.608**	0.030	0.750**	0.030
Social	0.673**	0.030	0.691***	0.039	0.250**	0.046	0.178**	0.062	0.691**	0.027	0.678**	0.035
Occupation	0.767**	0.029	0.778***	0.031	0.299**	0.048	0.127	0.068	0.727**	0.026	0.778**	0.028
Dignity	0.367**	0.040			0.059	0.041			0.421**	0.036		
Self-care			0.666***	0.035			0.240**	0.065			0.460**	0.051
Time & space			0.812***	0.026			0.198**	0.070			0.809**	0.026
Feel supported			0.540***	0.055			0.180**	0.054			0.260**	0.052
N	737		312		0.721**	.035	0.796**	.051	737		312	
χ^2	90.98**		30.67**		0.469**	.035	0.424**	.052	28.55**		14.61	
RMSEA	0.069	No	0.062	No	727		311		0.038	Yes	0.032	Yes
90% CI lower	0.055		0.032		13.43		33.27**		0.017		<0.001	
90% CI upper	0.084	No	0.092	No	0.036	Yes	0.121	No	0.057	Yes	0.072	Yes
SMSR	0.118	No	0.070	Yes	<0.001		0.083		0.022	Yes	0.024	Yes
CFI	0.788	No	0.890	No	0.064	Yes	0.163	No	0.980	Yes	0.994	Yes
TLI	0.703	No	0.835	No	0.013	Yes	0.038	Yes	0.969	Yes	0.991	Yes

*p<0.05, **p<0.01

¹ **Reflective:** Control, Social, Occupation, Dignity. **Formative:** Accommodation, Food & drink, Personal care, Personal safety.

² **Reflective:** Control, Social, Occupation, Time & space. **Formative:** Self-care, personal safety, Feeling supported.

All of the applied criteria were met for the ASCOT formative model, although the overall model fit (χ^2) was not significant. By contrast, the formative model was a poor fit for ASCOT-Carer. Again, only $\text{MSR} \leq 0.08$ was met. Finally, the mixed model was a good fit for ASCOT and ASCOT-Carer. Taken together, the findings indicate that the RMM is the worst fit for ASCOT and ASCOT-Carer. The FM or mixed models may apply to ASCOT. The mixed model was the best fit for ASCOT-Carer.

The network models are shown in Figure 7 and Figure 8. The **nodes** (A1-8, C1-7) represent the items in ASCOT and ASCOT-Carer. The **edges**, shown as lines between nodes, represent empirical correlation between nodes. Thicker lines represent a stronger correlation between items. For both the ASCOT analysis (Figure 7) and ASCOT-Carer (Figure 8), there are relevant edges between all nodes. All correlations were positive, shown by green lines.

Degree centrality, which is an indicator of the relative importance of each node that is defined by the number of connections incident to each node, ranged in the **ASCOT analysis** (Figure 7) from two (Dignity) to five connections (Accommodation). As expected, based on the theoretical concept of *higher order domains* and *basic domains*, there are relevant edges between A1 to A3 (higher order) and A5 to A7 (basic), with connection also to A4. The node that relates to the Dignity item has the lowest number of edges, to A3 (Occupation) and A7 (Accommodation), which aligns to the concept that it is distinct from the other nodes since it is the only one that relates to care delivery. The network for **ASCOT-Carer analysis** (Figure 8) also has

a degree centrality from two (Feeling supported) to five (Self-care). However, the structure does not align with the theoretical split between higher order domains (C1 to C3) and basic domains (C4 to C6) as for ASCOT.

The centrality indices for each node are reported in Table 3. These are indicators of the relative importance or influence of each node. Specifically, the indices represent a composite of the number and strength of connections (*strength*), the relationship with other nodes or its connectivity (*connectedness*) and the importance of a node in relation to the average pathway between other nodes (*betweenness*). For ASCOT (A1-8), the most influential nodes in ASCOT are Occupation and Personal cleanliness and comfort. Accommodation, Control and Personal safety are also indicated as influential by one of the strength, connectedness and betweenness indices, respectively. Since the strength index is least likely to be affected by sample size and is typically the most reliable of the three indices²⁹, we tentatively also highlight Accommodation as an influential node, alongside Occupation and Personal comfort and cleanliness. For ASCOT-Carer (C1-7), there are relevant edges between higher order domains (C1 to C3), but also C6 (Time and Space) to C1 (Control) and C3 (Occupation). The centrality indices indicate the most influential nodes are Time and space, Self-care, Social participation and Occupation.

Discussion

This study aimed to further understand the nature and internal structure of measures of SCRQoL for adults with social care needs (ASCOT) and their carers (ASCOT-Carer). In the development

Figure 7. Partial correlation network for ASCOT. Key: A1 (Control over daily life) A2 (Social participation) A3 (Occupation) A4 (Safety) A5 (Personal care) A6 (Food and drink) A7 (Accommodation) A8 (Dignity)

Figure 8. Partial correlation network for ASCOT-Carer. Key: C1 (Control over daily life) C2 (Social participation) C3 (Occupation) C4 (Safety) C5 (Self-care) C6 (Time and space to be yourself) C7 (Feeling supported and encouraged)

Table 3. Centrality indices.

	Dimension	Strength	Closeness	Betweenness
A1	Control over daily life	0.74	2.01	4
A2	Social participation	0.75	1.87	0
A3	Occupation	1.08	2.05	16
A4	Personal safety	0.55	1.93	6
A5	Personal comfort & cleanliness	0.90	1.94	6
A6	Food and drink	0.70	1.56	0
A7	Accommodation	0.94	1.87	2
A8	Dignity	0.31	1.39	0
C1	Control over daily life	0.86	2.19	0
C2	Social participation	0.83	2.50	4
C3	Occupation	0.94	2.47	2
C4	Personal safety	0.50	1.98	0
C5	Self-care	0.91	2.56	8
C6	Time and space	1.06	2.78	2
C7	Feeling support & encouraged	0.38	1.95	0

Note. The three (or four) most central nodes according to each index are reported in bold. Closeness values are multiplied by 100. Values may be compared within, not across, measures.

of the measures and psychometric testing of the original English language version or cultural adaptation and translations, authors have made different assumptions about whether the measure is formative^{1,2} or reflective^{15,16}. In some work, the issue of measurement models has been noted, and an ‘agnostic’ approach taken to analysis that does not commit to either model¹⁷. In this study, we have directly compared measurement models for the ASCOT and ASCOT-Carer as *reflective*, *formative* or *mixed*. This provides useful insights to inform the approach for future translation and cross-cultural adaptation of the ASCOT measures.

The findings indicate that the ASCOT and ASCOT-Carer do not empirically fit to the RMM. Based on this empirical finding, especially in combination with the literature on theoretical issues of applying RMMs to PBMs and/or health- or care-related QoL measures, like the ASCOT and ASCOT-Carer, we recommend that the ASCOT measures are not assumed to be reflective. By contrast, the analysis provides tentative empirical evidence of fit of the ASCOT to the formative or mixed model. The former is consistent with the development of the ASCOT, which argued that the attributes should be weakly correlated for its suitability as a PBM¹. The fit for the ASCOT-Carer to the formative model is, however, not supported by this analysis. Instead, the best fit is the mixed model. The current guidelines for the development of patient-reported outcome measures (PROMs) suggest that it is acceptable to apply reflective measurement methods (*i.e.* EFA) to examine structural characteristics of measures based on the mixed model⁸. This is relevant to cross-cultural adaptation and validation of translations, where EFA is recommended to establish cross-cultural validity⁸. Therefore, we recommend that future translations of the ASCOT measures may appropriately and usefully apply EFA to explore structural characteristics, especially if the translated version will be applied as a PBM in economic evaluations *and* as a measure of SCRQoL.

In addition to empirically testing the formative, reflective and mixed models, we also applied the network model to explore whether it could add further insights into the nature and structure of ASCOT and ASCOT-Carer. The findings align with the theoretical concept of ASCOT comprising higher order and basic domains that have stronger relationships, as has been found in previous analysis using EFA and correlations for ASCOT¹. The former relate to the domains of social participation, occupation and control, which may be conceptualized as aspects of care-related QoL/need beyond basic care-related QoL/needs, like accommodation, food and drink, safety and personal comfort and cleanliness. The analysis presented here for the ASCOT-Carer does not show the same division between basic/higher-order QoL/needs. However, there are key connections between the higher-order domains (*i.e.* social participation, occupation and control) and time and space to be yourself.

In the analysis presented here, the NM analysis provides useful insights into the nature and internal structure of the measures that add to the insights of established methods, like EFA, CFA or SEM. It does this without specifying the underlying measurement model and in a way that aligns more closely to the

complex relationships that are known to exist between aspects of health-related and social care-related QoL⁷. Specifically, the analysis for ASCOT indicates that Occupation (meaningful and enjoyable activity) and Personal comfort and cleanliness are the most influential aspects of QoL. Accommodation, control over daily life and personal safety are also key nodes. The most influential nodes for the ASCOT-Carer were Time and space, Self-care, Social participation and Occupation. Therefore, we propose that network analysis be used as a complementary approach in development and adaptation of ASCOT measures, alongside psychometric approaches, like EFA, to explore the internal structure and relationship between items.

Furthermore, the NM analysis presented here provides insights that may inform future qualitative research, as well as the application of the measures in care planning and assessment⁷. This is relevant as there has been interest in applying ASCOT in this way, in England and internationally^{13,14,40}. The most influential nodes for each measure and the key relationships between domains may be useful in informing needs assessment and care planning. Specifically, it may guide the conversation to focus on these aspects of QoL for service users (ASCOT) and carers (ASCOT-Carer) respectively, via the target of intervention (*e.g.* home care), to influence other aspects of QoL for additional benefit. In addition, it may guide research to understand whether and how specific social care interventions are effective. In previous studies, ASCOT measures have been used in qualitative interviews to identify how social care services impact on QoL of adults with care needs and carers^{1,41,42}. This research provides insight into which care supports QoL; however, less attention has been given to relationships between QoL domains. The nature of the relationship indicated by an edge in the NM may indicate a direct causal pathway or the common effect of a (latent) variable not included in model²⁹. These associations are indicative of causal relationships that require further investigation, drawing on qualitative evidence, that may then inform the design and evaluation of interventions⁶.

The study presented here has some limitations. The analysis was conducted on datasets from two studies in England. Further studies to replicate and confirm the findings, both in England and other countries with translated versions of the measure, would add further insight. Study 1 (IIASC) included a diverse sample of users of social care services with a range of needs. However, Study 2 (MOPED) included only a sub-group of carers, *i.e.* of people with dementia, who are known to have higher-level and specific needs compared to other carers. Modelling based on these data may not fully inform the structure of the ASCOT-Carer, when applied for use with carers more broadly, so replication with other groups of carers is important. Furthermore, the specification of the mixed models were limited in the choice of external measures (*i.e.* EQ-5D and overall QoL rating) by the variables available in the respective study datasets. Although the choice of these measures may be justified by previous studies that show the relationship between SCRQoL and the related constructs of health-related QoL and overall QoL^{1,12}, it may be that other external measures may be more suitable (*e.g.* carer-related quality of life or related measures based on the

capability approach, like the ICECAP-A⁴³ or ICECAP-O⁴⁴) and may lead to differences in the coefficients for structural paths and their significance. Indeed, we have not placed much emphasis on interpreting the structural paths, despite being of interest, for this reason.

Conclusions

The findings of this study indicate that the ASCOT and ASCOT-Carer are not adequately described as RMMs. The ASCOT fits best to either formative or mixed model. ASCOT-Carer fits best to the mixed model. These findings are relevant to cross-cultural adaptation and validation of translated versions of ASCOT and ASCOT-Carer. We recommend that future translations of the ASCOT measures may usefully and appropriately apply EFA to explore structural characteristics, especially if the translated version will be applied as a PBM in economic evaluations and as a measure of SCRQoL. Further investigation using datasets collected with different populations and in other contexts may usefully guide the approach and provide additional evidence of their internal structure. In addition to EFA, network analysis (based on the network model) may also provide useful insights into the relationships between items.

List of abbreviations

ASCOT	Adult Social Care Outcomes Toolkit
CFA	Confirmatory factor analysis
CFI	Comparative fit index
EFA	Exploratory factor analysis
FM	Formative measurement model
IIASC	Identifying the impact of adult social care study
LA	Local authority
MIMIC	Multiple indicators multiple causes model
MOPED	Measuring the outcomes of people with dementia and their carers study
NM	Network model
PBM	Preference-based measure
RMM	Reflective measurement model
RMSEA	Root mean square error of approximation
SCRQoL	Social care-related quality of life

SMSR	Standardized root mean square residual
TLI	Tucker-Lewis index
QoL	Quality of life

Declarations

Data availability

The datasets generated and/or analysed during the current study are not publicly available due to ethical considerations. Participants did not consent to their full data being shared outside of the research team. Reasonable requests for access to anonymised data will be considered. Please contact s.e.rand@kent.ac.uk.

Reporting guidelines

Not applicable.

Consent

IIASC: Ethical approval for the study was given by the Social Care Research Ethics Committee in England (12/IEC08/0049) with local research governance approval. Written informed consent to participate was obtained from all participants. **MOPED:** Ethical approval for the study was obtained from the Social Care Research Ethics Committee in England (19/IEC08/0057) with local research governance approval, where carers' organisations were involved in data collection. Approval to conduct the study in the NHS was granted from the Health Research Authority. Written informed consent to participate was obtained from all participants.

Author contributions

SR conceptualized the study, analyzed and interpreted the data and drafted the manuscript. A-MT and JM contributed to the concept. BS advised on the methodology. A-MT, JM and BS reviewed and edited the manuscript. All authors read and approved the final manuscript.

Acknowledgements

We would like to acknowledge and thank Dr. Birgit Trukeschitz and Professor Ismo Linnoosmaa for their contribution to the early development and discussion of the concept for this article. Recruitment to this study was supported through the National Institute for Health Research (NIHR) Clinical Research Network (CPMS: 44825) and Join Dementia Research, which is funded by the Department of Health and Social Care (DHSC) and delivered by the NIHR in partnership with Alzheimer Scotland, Alzheimer's Research UK and Alzheimer's Society.

References

- Netten A, Burge P, Malley J, *et al.*: **Outcomes of social care for adults: developing a preference-weighted measure.** *Health Technol Assess.* 2012; **16**(16): 1–166.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Rand SE, Malley JN, Netten AP, *et al.*: **Factor structure and construct validity of the Adult Social Care Outcomes Toolkit for Carers (ASCOT-Carer).** *Qual Life Res.* 2015; **24**(11): 2601–14.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Batchelder L, Malley J, Burge P, *et al.*: **Carer social care-related quality of life outcomes: estimating English preference weights for the Adult Social Care**

- Outcomes Toolkit for Carers (ASCOT-Carer).** Canterbury: Personal Social Services Research Unit, University of Kent, 2018.
4. Avila ML, Stinson J, Kiss A, et al.: **A critical review of scoring options for clinical measurement tools.** *BMC Res Notes*. BioMed Central, 2015; **8**: 612. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 5. Brazier J, Deverill M, Green C, et al.: **A review of the use of health status measures in economic evaluation.** *J Health Serv Res Policy*. 1999; **4**(3): 174–84. [PubMed Abstract](#) | [Publisher Full Text](#)
 6. Fayers PM, Hand DJ: **Factor analysis, causal indicators and quality of life.** *Qual Life Res*. 1997; **6**(2): 139–50. [PubMed Abstract](#) | [Publisher Full Text](#)
 7. Kossakowski JJ, Epskamp S, Kieffer JM, et al.: **The application of a network approach to Health-Related Quality of Life (HRQoL): introducing a new method for assessing HRQoL in healthy adults and cancer patients.** *Qual Life Res*. Springer International Publishing, 2016; **25**(4): 781–92. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 8. Mokkink LB, Prinsen CAC, Patrick D, et al.: **COSMIN Study Design checklist for Patient-reported outcome measurement instruments.** Amsterdam, 2019. [Reference Source](#)
 9. Mokkink LB, de Vet HCW, Prinsen CAC, et al.: **COSMIN Risk of Bias checklist for systematic reviews of Patient-Reported Outcome Measures.** *Qual Life Res*. Springer International Publishing, 2018; **27**(5): 1171–9. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 10. Forder JE, Caiels J: **Measuring the outcomes of long-term care.** *Soc Sci Med*. 2011; **73**(12): 1766–74. [PubMed Abstract](#) | [Publisher Full Text](#)
 11. van Leeuwen KM, Bosmans JE, Jansen APD, et al.: **Comparing Measurement Properties of the EQ-5D-3L, ICECAP-O, and ASCOT in Frail Older Adults.** *Value Heal*. 2015; **18**(1): 35–43. [PubMed Abstract](#) | [Publisher Full Text](#)
 12. Rand SE, Malley JN, Vadean F, et al.: **Measuring the outcomes of long-term care for unpaid carers: Comparing the ASCOT-Carer, Carer Experience Scale and EQ-5D-3 L.** *Health Qual Life Outcomes*. 2019; **17**(1): 184. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 13. Kettunen A: **Supporting Informal Carers: Testing Feasibility of the CES and the Carer SCT4 Measures as a Part of Care Practice.** 5th Int Conf Evidence-based Policy Long-term Care. 2018.
 14. Johnstone L, Page C: **Using Adult Social Care Outcomes Toolkit (ASCOT) in the assessment and review process.** *Res Policy Plan*. 2013; **30**: 179–92. [Reference Source](#)
 15. Trukeschitz B, Litschauer J, Hajji A, et al.: **Cross-cultural adaptation and construct validity of the German version of the Adult Social Care Outcomes Toolkit for service users (German ASCOT).** *Health Qual Life Outcomes*. 2020; **18**(1): 326. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 16. Trukeschitz B, Hajji A, Litschauer J, et al.: **Translation, cultural adaptation and construct validity of the German version of the Adult Social Care Outcomes Toolkit for informal Carers (German ASCOT-Carer).** *Qual Life Res*. Springer Science and Business Media Deutschland GmbH. 2020; **30**(3):905–920. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 17. Nguyen L, Linnosmaa I, Jokimäki H, et al.: **Social care-related outcomes in Finland. Construct validity and structural characteristics of the Finnish ASCOT measure with older home care users.** *Health Soc Care Community*. Blackwell Publishing Ltd. 2021; **29**(3): 712–28. [PubMed Abstract](#) | [Publisher Full Text](#)
 18. Feng YS, Jiang R, Kohlmann T, et al.: **Exploring the Internal Structure of the EQ-5D Using Non-Preference-Based Methods.** *Value Health*. Elsevier Ltd. 2019; **22**(5): 527–36. [PubMed Abstract](#) | [Publisher Full Text](#)
 19. Guyon H: **The Fallacy of the Theoretical Meaning of Formative Constructs.** *Front Psychol*. Frontiers Media S.A.; 2018; **9**: 179. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 20. Howell RD, Breivik E, Wilcox JB: **Reconsidering Formative Measurement.** *Psychol Methods*. 2007; **12**(2): 205–18. [PubMed Abstract](#) | [Publisher Full Text](#)
 21. Edwards JR: **The fallacy of formative measurement.** *Organ Res Methods*. 2011; **14**: 370–88. [Publisher Full Text](#)
 22. Edwards JR, Bagozzi RP: **On the nature and direction of relationships between constructs and measures.** *Psychol Methods*. 2000; **5**(2): 155–74. [PubMed Abstract](#) | [Publisher Full Text](#)
 23. Schmittmann VD, Cramer AOJ, Waldorp LJ, et al.: **Deconstructing the construct: A network perspective on psychological phenomena.** *New Ideas Psychol*. Elsevier Ltd. 2013; **31**: 43–53. [Reference Source](#)
 24. De Schryver M, Vindevogel S, Rasmussen AE, et al.: **Unpacking Constructs: A Network Approach for Studying War Exposure, Daily Stressors and Post-Traumatic Stress Disorder.** *Front Psychol*. Frontiers Media S.A.; 2015; **6**: 1896. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 25. Van Der Maas HLJ, Dolan CV, Grasman RPPP, et al.: **A dynamical model of general intelligence: The positive manifold of intelligence by mutualism.** *Psychol Rev*. 2006; **113**(4): 842–61. [PubMed Abstract](#) | [Publisher Full Text](#)
 26. Cramer AOJ, van der Sluis S, Noordhof A, et al.: **Dimensions of normal personality as networks in search of equilibrium: You can't like parties if you don't like people.** *Eur J Pers*. John Wiley & Sons. Ltd. 2012; **26**: 414–31. [Publisher Full Text](#)
 27. Eaton NR: **Latent variable and network models of comorbidity: toward an empirically derived nosology.** *Soc Psychiatry Psychiatr Epidemiol*. Springer Berlin Heidelberg. 2015; **50**(6): 845–9. [PubMed Abstract](#) | [Publisher Full Text](#)
 28. Borsboom D, Cramer AOJ: **Network Analysis: An Integrative Approach to the Structure of Psychopathology.** *Annu Rev Clin Psychol*. 2013; **9**: 91–121. [PubMed Abstract](#) | [Publisher Full Text](#)
 29. Hevey D: **Network analysis: A brief overview and tutorial.** *Heal Psychol Behav Med*. 2018; **6**(1): 301–28. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 30. Rand S, Malley J, Towers AM, et al.: **Validity and test-retest reliability of the self-completion adult social care outcomes toolkit (ASCOT-SCT4) with adults with long-term physical, sensory and mental health conditions in England.** *Health Qual Life Outcomes*. 2017; **15**(1): 163. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 31. Bollen KA: **Structural Equations with Latent Variables.** New York: Wiley; 1989. [Publisher Full Text](#)
 32. Malley JN, Towers AM, Netten AP, et al.: **An assessment of the construct validity of the ASCOT measure of social care-related quality of life with older people.** *Health Qual Life Outcomes*. 2012; **10**: 21. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 33. Netten A, Burge P, Malley J, et al.: **Outcomes of social care for adults: Developing a preference-weighted measure.** *Health Technol Assess*. 2012; **16**(16): 1–166. [PubMed Abstract](#) | [Publisher Full Text](#)
 34. van Hout B, Janssen MF, Feng YS, et al.: **Interim scoring for the EQ-5D-5L: Mapping the EQ-5D-5L to EQ-5D-3L value sets.** *Value Health*. Elsevier Inc.; 2012; **15**(5): 708–15. [PubMed Abstract](#) | [Publisher Full Text](#)
 35. Hu LT, Bentler PM: **Cutoff criteria for fit indexes in covariance structure analysis: Conventional criteria versus new alternatives.** *Struct Equ Model*. Taylor & Francis Group; 1999; **6**(1): 1–55. [Publisher Full Text](#)
 36. Friedman J, Hastie T, Tibshirani R: **Sparse inverse covariance estimation with the graphical lasso.** *Biostatistics*. 2008; **9**(3): 432–41. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 37. McCabe M, You E, Tatangelo G: **Hearing Their Voice: A Systematic Review of Dementia Family Caregivers' Needs.** *Gerontologist*. 2016; **56**(5): e70–88. [PubMed Abstract](#) | [Publisher Full Text](#)
 38. Larkin M, Henwood M, Milne A: **Older Carers and Carers of People with Dementia: Improving and Developing Effective Support.** *Social Policy and Society*. 2020; 1–15. [Publisher Full Text](#)
 39. Larkin M, Henwood M, Milne A: **Carer-related research and knowledge: Findings from a scoping review.** *Health Soc Care Community*. 2019; **27**(1): 55–67. [PubMed Abstract](#) | [Publisher Full Text](#)
 40. Towers A, Nelson K, Smith N, et al.: **Using ASCOT in Care Planning Conversations.** *Aust J Dement Care*. 2018.
 41. Rand S, Vadean F, Forder J: **The impact of social care services on carers' quality of life.** *Int J Care Caring*. Policy Press; 2020; **4**(2): 235–59. [Publisher Full Text](#)
 42. Rand S, Malley J: **Carers' quality of life and experiences of adult social care support in England.** *Health Soc Care Community*. 2014; **22**(4): 375–85. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 43. Al-Janabi H, Flynn TN, Coast J: **Development of a self-report measure of capability wellbeing for adults: the ICECAP-A.** *Qual Life Res*. Springer Netherlands; 2012; **21**(1): 167–76. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 44. Grewal I, Lewis J, Flynn T, et al.: **Developing attributes for a generic quality of life measure for older people: Preferences or capabilities?** *Soc Sci Med*. Pergamon; 2006; **62**(8): 1891–901. [PubMed Abstract](#) | [Publisher Full Text](#)

Open Peer Review

Current Peer Review Status: ? ?

Version 1

Reviewer Report 21 October 2022

<https://doi.org/10.3310/nihropenres.14375.r28794>

© 2022 Sufriyana H. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Herdiantri Sufriyana

¹ Graduate Institute of Biomedical Informatics, College of Medical Science and Technology, Taipei Medical University, Taipei, Taiwan

² Department of Medical Physiology, Faculty of Medicine, Universitas Nahdlatul Ulama Surabaya, Surabaya, Indonesia

General comment

This study compared fitting of three methods (i.e., reflective, formative, and mixed) to develop a model for measuring social care-related quality of life (ASCOT) as the construct using several observables. The models were developed for measuring the construct in a care recipient (i.e., ASCOT) or giver (i.e., ASCOT-Carer or ASCOT-C). Subsequently, methodologically validation of these models was conducted by network analysis of the observables. This study is important for future translation and cross-cultural adaptation of the ASCOT measures. Social observables may interact differently across cultures. Structural relationships of the measurement model in the original culture should be described to identify such difference. Nevertheless, several major issues were identified in this study, which may make this study inconclusive yet (please kindly see major comments). Note, reviewer particularly commented on the methodological element of this manuscript, leaving the domain-field element uncommented, due to the limitation of the reviewer's expertise.

Major comments

1. Why were no uncertainty intervals for SMSR and CFI/TLI? To differ if a model indicated a good fit using the criteria from a finding which simply happened by chance (spurious correlation or sampling error), it is important to include the intervals, such as 95% confidence interval. This is possible to be inferred by resampling.
2. How did this study deal with multiple-testing effect? This study only assessed the significant association by p-values, which resulted in many significant findings. Please kindly consider to correct the p-values for multiple-testing effect. Also consider to take effect size into account when identifying the significant association.

3. How was strength, closeness, and betweenness computed? Please briefly explain how to get the numbers. It is also unclear how to deal with these three dimensions to achieve the analytical conclusion.

Minor comments

1. Please use terms consistently, e.g.: (1) closeness vs. connectedness which was interchangeably used in the main text and Table 3; (2) ASCOT-Care vs. ASCOT-C; and (3) other terms not limited to these examples.
2. Please kindly consider to classify basic and higher-order domains with sublabels in Tables 2 and 3. Also add sublabels for ASCOT and ASCOT-C in Table 3. It is easier than remembering the alphanumeric codes of the observables.
3. The Introduction and Discussion seem to be written divergently, which makes it difficult to capture the central message. It is still unclear how this methodological comparison plus its network analysis would eventually help for future translation and cross-cultural adaptation of the ASCOT measures. This probably answers the “so-what” of this study.
4. Please kindly consider to share the analytical codes, particularly those by R.

Is the work clearly and accurately presented and does it cite the current literature?

Yes

Is the study design appropriate and is the work technically sound?

Yes

Are sufficient details of methods and analysis provided to allow replication by others?

Yes

If applicable, is the statistical analysis and its interpretation appropriate?

Partly

Are all the source data underlying the results available to ensure full reproducibility?

Partly

Are the conclusions drawn adequately supported by the results?

Partly

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: Causal inference from observational data, Statistical and computational machine learning, Pathway (network) analysis.

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard, however I have significant reservations, as outlined above.

Author Response 09 Mar 2023

Stacey Rand

Response. Thank you for your helpful and constructive comments. Our response is below.

Major comments

1. Table 2 now includes the 95% CI estimates for the RMSEA, SMSR and CFI/TFI. The text has been edited to clarify that we apply a goodness-of-fit criteria for the RMSEA based on the 90% confidence interval (see Methods, p.9, following Hu & Bentler, 1999), so we also report the 90% CI for the RMSEA.

In adding the 95% CI estimates, we noticed and corrected two errors in Table 2. (These are shown in track changes for ease of reference). These do not substantively change the results, but we have also highlighted the changes in the report of the results (p.13).

i. The goodness-of-fit statistics in the MIMIC formative models for ASCOT and ASCOT-Carer were incorrect, due to a formatting error in the previous version. This has led to values showing on the wrong row of the Table, with the omission of standard coefficients/SE for the QoL measures (EQ-5D and QoL item) and the CFI/TLI. These are now correctly presented, along with the 95% CI estimates.

ii. We realised that there was an inconsistency across the models. The formative models were reported using weighted least squares; the other models using maximum likelihood. For consistency, all analyses are now reported using maximum likelihood, which is what we had originally intended. We have also corrected the description in the Methods section (p.7). The criteria proposed by Hu & Bentler, 1999 – based on ml estimation – are then applied. This does not substantively change the key findings (i.e. that RMM is not a good fit for ASCOT or ASCOT-Carer), even if some of the criteria that were not previously met are now met (i.e. $\text{SMSR} \geq .08$ for ASCOT and $\text{CFI/TLI} \geq .95$ for ASCOT-Carer).

2. The analyses (CFA, MIMIC and network analysis) do not apply any correction for multiple comparisons. The rationale is that the study is designed to understand the structural characteristics of the ASCOT and ASCOT-Carer, against models based on reflective, formative and mixed measurement models. The focus is not on specific factor loadings or standardised coefficients for each item/variable; rather, we are seek to test the fit against the models. This is reflected in the approach in the paper (i.e. a focus on overall model fit, goodness-of-fit statistics).

3. The centrality indices (strength, closeness and betweenness) were calculated using R (see attached syntax file). These indices are described in the Methods (p.9):

“Centrality indices were also calculated and reported for node strength, closeness and betweenness. The strength index is a composite measure of both the number and strength of connections to a node. The closeness index represents the relationship between one node and the other nodes through its indirect connections (i.e. its connectedness or connectivity). A high closeness index indicates that the node is affected quickly by changes to the other nodes in the network. The betweenness index indicates the importance of a

node in relation to the average pathway between other nodes.”

Reference: Hevey D: Network analysis: A brief overview and tutorial. *Heal Psychol Behav Med.* 2018;6(1):301–28. 34040834 10.1080/21642850.2018.1521283 8114409

The Results section (p.15) has been edited to improve the clarity of the reported partial correlation networks and their interpretation (Figure 7, 8). The centrality indices are reported in Table 3 and described on pp.15-16.

Minor comments

1. The manuscript has been reviewed and edited for consistency of terminology - specifically, ASCOT-C has been edited to ASCOT-Carer; connectedness to closeness, when referring to the centrality indices.
2. Tables 2 and 3 have been edited to state which domains relate to ASCOT and/or to ASCOT-Carer, as well as the sublabels of *basic* or *higher order* domains. In Table 3, there are sublabels for each of the domains, alongside the alphanumeric codes.
3. The introduction (pp.3-5) has been edited to further clarify the link between this study and its application to the psychometric evaluation of translated and cross-cultural adapted PBMs.

This is also explained in the next paragraph of the Introduction (p.4), as below, as well as the final paragraph of the Introduction (p.5).

“Inconsistent approaches to psychometric evaluation of the same measure have given rise to discussion of the correct approach to the translation and cross-cultural validity or adaptation of the measures for new contexts or populations 15– 17.”

This is then restated in the Discussion, paragraph 1 (p.17) and further explored through the Discussion. Some minor edits have been made to the conclusion to, again, clarify how these findings are relevant to translation and cross-cultural adaptation (i.e. they inform our recommendation of the analytic approach to psychometric evaluation of translated and adapted measures).

4. The STATA and R syntax are provided as an appendix.

Competing Interests: None.

Reviewer Report 07 March 2022

<https://doi.org/10.3310/nihropenres.14375.r28513>

© 2022 Feng Y. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

You-Shan Feng

Institute for Clinical Epidemiology and Applied Biostatistics, Medical University of Tübingen, Tübingen, Germany

The authors present a very interesting paper looking into which model has the best fit with ASCOT data. The study contributes important work to determine how best the items of the ASCOT can be presented and perhaps scored use outside of CUA. The paper is very clearly presented and well written. I also appreciate the detail and presentation of the figures and tables. I have just a few points:

1. I wonder why only 1-factor models were used for the reflective model. Especially as the RMM is found to be the worst fit for the data, it might be prudent to investigate whether an alternative RMM could result in a better fit. Perhaps a 2- or 3-factor model, or a higher order model, would results in worse fit, but this should be discussed in the paper.
2. Why was the preference-based EQ-5D scoring system used instead of a level sum score or using the items as is? Similarly, was the EQ-VAS considered?
3. Given the spread of responses for ASCOT in Study 1, I am very surprised by how low the mean EQ-5D score is: 0.27. The low EQ-5D score should be mentioned in the discussion section as well: is this an expected mean score for this population? How does this and ASCOT responses compare to other patient populations?
4. A terminology issue: while it is perfectly appropriate to use the term “mixed measurement model” to describe models which include both RMM and FM, the term “mixed model” can be confusing as the terminology is also applied to hierarchal regression methods. I advise authors to consider revising the term “mixed model” for clarity.

From my point of view, the manuscript is ready for indexing once these few points of concern are addressed.

Is the work clearly and accurately presented and does it cite the current literature?

Yes

Is the study design appropriate and is the work technically sound?

Yes

Are sufficient details of methods and analysis provided to allow replication by others?

Yes

If applicable, is the statistical analysis and its interpretation appropriate?

Yes

Are all the source data underlying the results available to ensure full reproducibility?

Yes

Are the conclusions drawn adequately supported by the results?

Yes

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: Clinical Epidemiology, Biostatistics, Psychometrics, Instrument development, Questionnaire scoring, EuroQol Instruments, Self-rated health, Quality of Life Questionnaires

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard, however I have significant reservations, as outlined above.

Author Response 09 Mar 2023

Stacey Rand

Response. Thank you for your helpful and constructive comments. Our response is below.

1. We considered one-factor models based on previous studies using exploratory factor analysis, which have found the ASCOT and ASCOT-Carer to fit to a one-factor model (Netten *et al*, 2012; Rand *et al*, 2019). The manuscript has been edited (p.7.) to explain this. In addition, EFA for the ASCOT and ASCOT-Carer were calculated with the study datasets (IIASC and MOPED, respectively). In both cases, the EFA indicated a one-factor solution.
2. The preference-based EQ-5D scoring system was applied based on current guidance on how to derive an index score from the five EQ-5D-3L/5L item scores for UK studies. Previous studies have found that the ASCOT/-Carer are moderately positively correlated with the EQ-5D-3L, whereas not all items were significantly associated with ASCOT SCRQoL (e.g. the association between ASCOT-Carer and EQ-5D self-care was not significant (Rand *et al*, 2019)). This informed our decision to focus on the EQ-5D index score. The EQ-VAS was not considered in the analysis. It was only collected in the MOPED study, not the IIASC study. To reflect this, the limitations section has been amended to also refer to the EQ-VAS.
3. The low EQ-5D score reflects the IIASC study aim, design and sample, i.e., it was a study of adults, who use community-based adult social care services (e.g., homecare to help with washing, dressing, eating) due to support needs related to long-term health conditions or disability. ASCOT is a measure of adult social care outcomes designed for completion by adults using adult social care services (e.g. home care). Therefore, studies that apply ASCOT (like IIASC) are likely to have samples with lower EQ-5D scores than general population. The introduction (p.3), methods (p.5) and results (p.10) have been edited to make this clearer.
4. The term 'mixed model' has been revised to 'mixed measurement model' throughout.

Competing Interests: None.